

CERENIA® with Certainty

Your Guide to Accurate Dosing




Cerenia Injectable Solution contains maropitant citrate 10mg/mL. It is indicated for:

- The prevention of vomiting in dogs and cats, except that induced by motion sickness.
- The treatment of acute vomiting in cats over 16 weeks of age, dogs and puppies 8 weeks of age or older, in combination with other veterinary and supportive measures.
- Aiding in the prevention and treatment of nausea.
- Prevention of perioperative nausea and vomiting and improvement in recovery from general anaesthesia after use of the μ -opiate receptor agonist morphine.




TOP TIPS FOR CERENIA USE

Tip #1: For prevention of nausea and vomiting, Cerenia Injectable should be administered more than 1 hour in advance. The duration of effect is approximately 24 hours, and therefore treatment can be given the night before administration of an agent that may cause nausea or emesis e.g. chemotherapy.



Dosing Chart for Dogs

Bodyweight (kg)	Dose Volume (mL)
1	0.1
2	0.2
3	0.3
4	0.4
5	0.5
6	0.6
7	0.7
8	0.8
9	0.9
10	1.0
12	1.2
14	1.4
16	1.6
18	1.8
20	2.0
25	2.5
30	3.0
35	3.5
40	4.0
45	4.5
50	5.0
55	5.5
60	6.0



Dosing Chart for Cats

Bodyweight (kg)	Dose Volume (mL)
1	0.1
1.5	0.15
2	0.2
2.5	0.25
3	0.3
3.5	0.35
4	0.4
4.5	0.45
5	0.5
5.5	0.55
6	0.6
6.5	0.65
7.0	0.7
7.5	0.75

Cerenia Injectable Solution should be administered **subcutaneously or intravenously**, once daily at a dose of 1 mg per kg bodyweight (1 mL per 10 kg bodyweight). Treatment may be repeated for up to five consecutive days.



TOP TIPS FOR CERENIA USE

Tip #2: If not injecting intravenously, injecting the product at refrigerated temperatures may reduce pain at the time of subcutaneous injection.