




Dosing Chart: Tablets

For prevention of vomiting due to motion sickness

For dogs 16 weeks and older, administer Cerenia tablets orally at 8 mg/kg once daily for up to two consecutive days. Administer at least one hour prior to travel. Do not redose if vomiting is observed. The duration of effect is at least 12 hours, which may allow administration the night before early morning travel.



General Instructions:

A light meal or snack approximately one hour prior to administration will reduce the incidence of post-treatment vomiting. Do not wrap or encapsulate in food as this may delay dissolution of the tablet and consequently the onset of effect.

Body weight (kg)	Number of Tablets			
	16 mg	24 mg	60 mg	160 mg
0.8 - 1.0	½			
1.1 - 1.5		½		
1.6 - 2.0	1			
2.1 - 3.0		1		
3.1 - 4.0	2			
4.1 - 5.0	2½			
5.1 - 6.0		2		
6.1 - 7.5			1	
7.6 - 10.0				½
10.1 - 11.3			1½	
11.4 - 15.0			2	
15.1 - 20.0				1
20.1 - 22.5			3	
22.6 - 30.0				1½
30.1 - 40.0				2
40.1 - 50.0				2½
50.1 - 60.0				3

For prevention of nausea and acute vomiting

Body weight (kg)	Number of Tablets		
	16 mg	24 mg	60 mg
3.0 - 4.0*	½		
4.1 - 8.0	1		
8.1 - 12.0		1	
12.1 - 24.0		2	
24.1 - 30.0			1
30.1 - 60.0			2

* The correct dose for dogs of less than 3kg cannot be accurately achieved with tablets.

For dogs and puppies 8 weeks and older, administer Cerenia tablets orally at 2 mg/kg once daily for up to 14 consecutive days. For prevention of nausea and vomiting, administer more than one hour in advance. The duration of effect is approximately 24 hours, therefore treatment may be given the night before administration of a potential emetic, e.g. chemotherapy. Cerenia tablets may be used interchangeably with Cerenia injectable solution for once daily dosing for the prevention of acute vomiting.